



Overview of studies on ecosystem services in riparian zones: a systematic review

Panorama dos estudos sobre serviços ecossistêmicos em zonas ripárias: uma revisão sistemática

Rachel Bardy Prado^{1*} , Gustavo Menezes Silva Damasceno²  and Fabiana de Gois Aquino³ 

¹Embrapa Solos, Rua Jardim Botânico, 1024, Jardim Botânico, CEP 22460-000, Rio de Janeiro, RJ, Brasil

²Escola Nacional de Saúde Pública, Fundação Oswaldo Cruz, Rua Leopoldo Bulhões, 1480, Mangueiras, CEP 21041-210, Rio de Janeiro, RJ, Brasil

³Embrapa Cerrados, BR 020, Km 18, Planaltina, CEP 73310-970, Brasília, DF, Brasil

*e-mail: rachel.prado@embrapa.br

Cite as: Prado, R.B., Damasceno, G.M.S. and Aquino, F.G. Overview of studies on ecosystem services in riparian zones: a systematic review. *Acta Limnologica Brasiliensia*, 2022, vol. 34, e19.

Abstract: Aim: Riparian zones are highly complex ecosystems, located on the banks of water bodies, with a fundamental role in maintaining biodiversity and ecosystem services (ES). This study aimed to systematize the knowledge about studies on ES in riparian zones, emphasizing methodological aspects and pointing out gaps and opportunities to reinforce their importance. **Methods:** The study was carried out based on literature review data over a period of 21 years (2000-2020), using Scopus and Web of Science databases. In the first stage, aspects of bibliometrics were analyzed, as well as the countries that published the most on the subject. In the second stage, the methodological aspects were analyzed (with emphasis on the integrated analysis of multiple ES, which looked at the landscape, adopted multiscale or carried out economic valuation). **Results:** From 2000 to 2020, 6,969 publications were obtained from Scopus and 16,498 from Web of Science, applying the search terms riparian buffer or similar and 371 publications were obtained from Scopus and 1,512 from Web of Science applying ecosystem service and riparian zones or similar terms, with the USA being the country that most published about ES in riparian zones. From a total of 219 publications selected, the ES category most studied in riparian zones was Regulation (65%), followed by Support (16%), Provision (8%) and Cultural (2%). Publications that studied three or more ES corresponded to 9% of the analyzed publications. Approximately 10% of publications used methodological approaches with multiple ESs in an integrated way. Less than 10% of publications focused on economic valuation. Gaps and opportunities were identified concerning the relevance and methods for evaluating and valuing ESs in riparian zones. **Conclusions:** Few studies used methodological approaches integrating different ES. That calls attention to the need to carry out more studies that analyze ES in riparian zones using an integrated and multiscale approach because that is how the components of the ecosystem interact and provide joint responses that may assist in decision making.

Keywords: integration; landscape; economic valuation; gaps; opportunities.

Resumo: Objetivo: As zonas ripárias são ecossistemas de alta complexidade, localizados às margens de corpos hídricos, com papel fundamental na manutenção da biodiversidade e dos serviços ecossistêmicos (SE). Este estudo teve como objetivo sistematizar o conhecimento sobre os SE em zonas ripárias, visando identificar os aspectos metodológicos e apontar lacunas e oportunidades para reforçar a sua importância. **Métodos:** O estudo foi realizado com base em dados da revisão da literatura



durante um período de 21 anos (2000-2020), usando as bases Scopus e Web of Science. Na primeira etapa foram analisados aspectos da bibliometria, bem como os países que mais publicaram no tema. Na segunda etapa foram analisados os aspectos metodológicos (com ênfase na análise integrada de múltiplos SE, que olharam para a paisagem, adotaram multiescalas ou realizaram valoração econômica). **Resultados:** De 2000 a 2020, 6.969 publicações foram obtidas da base Scopus e 16.498 da Web of Science, aplicando os termos de busca zonas ripárias ou similares e 371 publicações foram obtidas da base Scopus e 1.512 da Web of Science aplicando os termos serviços ecossistêmicos e zonas ripárias ou termos similares, sendo os EUA o país que mais publicou sobre ES em zonas ripárias no período analisado. De um total de 219 publicações, a categoria de SE mais estudada em zonas ripárias foi a de Regulação (65%), seguida de Suporte (16%), Provisão (8%) e Cultural (2%). As publicações que estudaram três ou mais SE corresponderam à 9% das publicações analisadas. Aproximadamente 10% das publicações utilizaram abordagens metodológicas com múltiplos SE de forma integrada. Menos de 10% das publicações focaram em valoração econômica. Lacunas e oportunidades foram identificados em relação à relevância e métodos para avaliação e valoração dos SE em zonas ripárias. **Conclusões:** Notou-se que poucos estudos utilizaram abordagens metodológicas integrando diferentes SE, utilizando multiescalas e fazendo a valoração econômica dos SE, o que chama a atenção para a necessidade de realizar mais estudos que analisem a oferta de ES por zonas ripárias de forma integrada, pois é de forma complexa e em diferentes escalas que os componentes do ecossistema interagem e fornecem respostas conjuntas capazes de auxiliar na tomada de decisões.

Palavras-chave: integração; paisagem; valoração econômica; lacunas; oportunidades.

1. Introduction

The riparian zone is the interface between the aquatic and terrestrial ecosystems, where the natural vegetation is usually present along the rivers, streams, lakes, and dams (Gregory et al., 1991; Kobiyama, 2003; Riis et al., 2020). The ecosystem comprises the interaction system of living things together with their non-living habitat, producing circulation, transformation, and accumulation of energy and matter (Tansley, 1935). As ecotones, riparian zones encompass sharp gradients of environmental factors, ecological processes, and plant communities. Riparian zones are not easily delineated but are comprised of mosaics of landforms, communities, and environments within the larger landscape (Gregory et al., 1991).

Riparian zones and their components interact generating important functions that include, in general, bank stabilization, provision of living and dead organic matter, habitat for both aquatic and terrestrial biota, the capture of sediment, retention and processing of nutrients, and moderation of the extreme temperatures by providing shade. These functions are the result of natural processes, which comprise the complex interactions between biotic and abiotic environments (De Groot et al., 2002). The ecosystem functions can be translated into ecosystem service (ES) as they trigger a series of benefits to human wellbeing (Costanza et al., 1997, 2017; Power, 2010). A single ES can be the product of two or more ecosystem functions, or a single ecosystem function can generate more than one ES. The ecosystem services (ES) provided by

riparian zones are closely related to water and some authors call them hydrologic ecosystem services (e.g. Brauman et al., 2007). Hydrologic ES are the benefits received by human from water ecosystems (Brauman et al., 2007) and Tabacchi et al. (2000) presented some impacts of riparian vegetation on hydrological processes. Although riparian zones have a multifunctional and fundamental role in the provision of ES they are strongly impacted due to anthropogenic pressures, especially during the urbanization process, in which the riverbanks were prioritized for occupation and had their riparian vegetation eliminated or reduced (Tiegs et al., 2019; Cao & Natuhara, 2020). Sweeney et al. (2004) and Sweeney & Newbold (2014) discussed the relationship between riparian deforestation, loss of stream and ecosystem services provision. Lind et al. (2019) pointed out guidelines to protect ecosystem functions and biodiversity in riparian zones present in agricultural landscapes, and Riis et al. (2020) identified, described, and ranked ES.

Riparian zones and riparian vegetation have been studied from many perspectives that cover multiple scientific and applied disciplines such as hydrology, biology, geography, remote sensing, management, and restoration (Riis et al., 2020). Hence, knowledge on this subject is distributed among a wide range of fields (Dufour et al., 2019).

Quantitative assessment of ecosystem services, functions, and flows is essential to maintain the ecological functions that riparian areas provide (Fu et al., 2016). However, it is observed that ES assessments in RZ are dispersed and there are gaps related to integrated methodological

approaches, focusing on landscape, multiscale, and economic valuation aspects, since riparian zones are ecosystems with high interaction and complexity. Perspectives based on isolated components are ecologically incomplete (Gregory et al., 1991) and often have very limited performance in spatial terms, not considering the landscape scale and not using economic valuation methods. Perspectives based on isolated components are unable to assess the trade-offs between the provision of different ES. In this context, we sought to identify in the international literature which the ecosystem services and methodologies have been applied for their evaluation and economic valuation, highlighting gaps and opportunities aiming to analyze multiple ecosystem services in an integrated manner in riparian zones.

2. Material and Methods

Several terms can mean the area or vegetation that occupies the margins of water bodies. The terms most found in the publications on ES in these areas were: riparian zones followed by riparian vegetation/forest, riparian buffer, and riparian areas. Therefore, these were the terms used in the systematic literature search and the term “riparian zones” was adopted and used throughout the text in this study. Some concepts treat the riparian zones from the point of view of the physical space occupied by the vegetation or the area located adjacent to the water body (Torres et al., 1992;

Bren, 1993; NRCS, 1997; Souza, 1999; Rodrigues, 2000; Selles, 2001; McKergow et al., 2003; Webb & Erskine, 2003) while others explicitly consider the interaction between terrestrial and aquatic ecosystems (Salvador, 1987; Gregory & Ashkenas, 1990; Gregory et al., 1991; Kobiyama, 2003; Vitalli et al., 2009; Attanasio et al., 2012; Aguiar Junior & Parron, 2015; Riis et al., 2020). Different definitions both based on geomorphology and biological interactions were equally considered, without applying judgment on which term would be more appropriate.

The present study was conducted based on a literature review, to answer two key questions:

- How has the evaluation of ecosystem services in riparian zones occurred in the last two decades (methodological aspects - main ES and indicators)?
- Do the applied methodologies for ES evaluation in riparian zones integrate different ecosystem services, scales and carry out their economic valuation?

Figure 1 presents the main steps of the study.

2.1. Step 1: Bibliographic survey

A literature review was carried out applying a quantitative systematic method (Denyer & Tranfield, 2009). Two databases of international publications (Scopus and Web of Science) were




Figure 1. Main steps of this study.

used, from 2000 to 2020, considering papers published in journals and conference proceedings. The search terms were applied in three steps as shown in Table 1 and the searches were performed using the fields: title, abstract and keywords of the manuscripts. The purpose of the first search was to get an idea of the trend of publications on the theme of “riparian zones” and similar terms (called of all terms), to compare with the results of the second search that included the term ecosystem service. In relation to the countries, the affiliation of the authors was considered, in order to know which countries are studying and investing more in the central theme of the article. It was not the scope of this article to analyze the study area of each article. The third search aimed to identify the publications that studied the selected ecosystem services, adopting the criteria presented below and in Table 1, in riparian zones or similar terms (called of all terms). In this step, publications found both in the Scopus and Web of Science databases were counted only once. In addition, in this step publications that studied three or more ecosystem services were grouped into another category (called several ES).

As the types of ES are many, as well as their classification, we chose to organize the ES according to the classification of Costanza et al. (1997) and Soman et al. (2007). While Costanza et al. (1997) considered “water supply” as a regulating ES, in the present study we chose to consider it in the provision ES category, as used by Grizzetti et al. (2016), which is a synonym for “water provisioning for drinking”.

We also added carbon stock/sequestration/carbon storage and nutrient retention ES, due to its importance in riparian zones, although it was not contemplated by Costanza et al. (1997), Soman et al. (2007), and Grizzetti et al. (2016). Searches were performed on titles, abstracts, and keywords of publications in both databases Scopus and Web of Science. Due to the proximity and function shading in the studies surveyed, some ecosystem services were grouped, such as: carbon storage, carbon sequestration, gas regulation (greenhouse gases), and climate regulation; erosion control and sediment retention; nutrient cycling and soil formation; nutrient retention and sewage treatment; nursery and refugia; food production and raw material and recreation and aesthetic. Table 2 shows the ES organized by category used in the present study. The main ecosystem functions and flows addressed in these publications were also identified at this step.

2.2. Step 2: Analysis of the methodological aspects applied to analyze ES in riparian zones

In this step, only the methodological aspects of publications that focused on ES in riparian zones (and similar terms) were analyzed (resulting from the third search). The analysis carried out from the publications that studied ecosystem services, according to the cut made in the present study, were analyzed regarding the most used indicators, integration methods of different ecosystem services, if they approached landscape and multiscale and if economic valuation methodologies were applied.

Table 1. Terms and methodology of the search in the literature used.

Search 1		Search 2		Search 3	
Terms	1 st terms	2 nd term	1 st terms	2 nd term	
OR “riparian buffer” “riparian vegetation” “riparian zone” “riparian area” “riparian forest”	OR “riparian buffer” “riparian vegetation” “riparian zone” “riparian area” “riparian forest”	AND “ecosystem service”	Search performed separately for each ES “biological control” “pollination”	AND “ecosystem service”	“erosion control” OR “sediment retention” “nutrient retention” OR “sewage treatment” “carbon stock” OR “sequestration” OR “gas regulation” OR “climate regulation” “water supply” “water regulation” “nutrient cycling” OR “soil formation” “food production” OR “raw material” “nursery” OR “refugia” “recreation” OR “aesthetic”

This analysis was not quantitative but sought to bring out an overview of what has been applied. The publications obtained were read in full to obtain the information necessary to carry out the analysis that was intended. The discussion of results was based on gaps and opportunities for research and decision-making support linked to the topic addressed.

3. Results

3.1. Bibliometrics

3.1.1. Riparian zones and ecosystem services in riparian zones terms (2000-2020)

From 2000 to 2020, 6,969 publications were obtained from Scopus and 16,498 from Web of Science, applying the search terms: riparian buffer or riparian vegetation or riparian zone or riparian areas or riparian forest (all terms) (Figure 2a). 371 publications were obtained from Scopus and 1,512 from Web of Science applying ecosystem service and riparian zones and similar terms (Figure 2b). It is observed that the growth in the number of publications in the different databases follows the same trend of publications that address

riparian zones and similar terms, without using the ES approach. The United States was the country that has published most about all terms related to riparian zones (45%), followed by Brazil (10%), Canada (8%), China (7%), Australia (7%) and others (23%). The USA also has published more when the focus was on ES in riparian zones (55%), followed by Canada (8%), Brazil (6%), German (5%), United Kingdom (5%), Australia (5%) and others (16%). The United States has invested heavily in studies on riparian zones in recent years to mitigate and remedy their degradation, because between 1780 and 1980, 53% of these areas were destroyed in the 48 states of the United States (Dahl, 1990; Mitsch & Gosselink, 2000). The high investments in science and research turned the United States into one of the countries that most publishes scientific articles in the world according to SCImago (2020).

3.1.2. Ecosystem services most studied in riparian zones

A total of 219 publications were obtained by applying the terms of the third search. The ES category most studied in the riparian zones was

Table 2. Ecosystem services provided by riparian zones by categories.

Ecosystem services categories	Ecosystem services
Provisioning	Water supply, food production, raw materials, and genetic resources
Regulating	Biological control, erosion control, sediment retention, nutrient retention, disturbance regulation, climate regulation, gas regulation, water regulation, pollination, waste treatment, carbon stock or sequestration
Support	Nutrient cycling, soil formation, nursery and refugia
Cultural	Aesthetic and recreation

Source: Modified from Costanza et al. (1997), Soman et al. (2007) and Grizzetti et al. (2016).




Figure 2. (a) Number of publications annually from 2000 to 2020 applying all terms. (b) Number of publications annually from 2000 to 2020 applying all terms and ecosystem services (Scopus and Web of Science databases).

Regulating (65%), followed by Support (16%), Provisioning (8%) and Cultural (2%). Publications that studied three or more ecosystem services were grouped into another category (called several ES), equivalent to 9% of the analyzed publications (Figure 3).

In the Regulating category, the ES studied were carbon stock or sequestration, gas regulation (greenhouse gases) and climate regulation (26%), biological control (17%), pollination (14%), erosion control and sediment retention (14%), water regulation (13%), nutrient retention and sewage treatment (11%), and disturbance regulation (5%). In the Provisioning category, the ES studied were water supply (76%) and food production and raw material (24%). Genetic resources ES was not found in the analyzed publications and if we included the term “agriculture” in the search, the number of publications related to food production and raw material would greatly increase, but this was not the focus of the study. The Support category included the ES of nutrient cycling and soil formation (56%) and nursery, refugia (44%). In the Cultural category, the ES studied were Recreation and aesthetic (Figure 3).

3.2. Ecosystem services in riparian zones: analysis of the methodological aspects

3.2.1. Ecosystem services indicators in riparian zones

Approximately 25% of the publications resulting from the third search that studied ES in

RZ (219 publications) and stood out for using the Millenium Ecosystem Assessment (MEA, 2005) as a reference. Table 3 shows the ES and example of indicators used in the analyzed publications.

3.2.2. Methodologies of integrated analysis of ecosystem services in riparian zones

Approximately 9% of the publications, resulting from the third search that studied ES in RZ (219 publications), studied three or more ES, and it was in these cases that methodologies of integrated analysis of ecosystem services were applied. The list with the main methods of integrated analysis of ES used (models and indexes), the objective of the analysis, and some references related can be found in Table 4.

3.2.3. Ecosystem services zone at landscape or spatial multiscale in riparian

Approximately 15% of the publications resulting from the third search that studied ES in RZ (219 publications), used landscape scales or spatial multiscale. The most of them aimed to evaluate the impact of human activities on the natural landscape in riparian areas and understand how these activities can jeopardize the provision of ES.

3.2.4. Economic valuation of ecosystem services in riparian zones

Less than 10% of the publications resulting from the third search that studied ES in RZ (219 publications), focused on the economic




Figure 3. Number of occurrences found for ecosystem service (ES) in riparian zones, according to the classification by Costanza et al. (1997), Soman et al. (2007) and Grizzetti et al. (2016).

Table 3. Main indicators used to assess ecosystem services in riparian zones.

ES category	Ecosystem services	Example of indicators
Provisioning	Water supply	Infiltration of water in the soil, slope, and water entering the system
	Raw material	Raw material collected, wood biomass
	Food production	Arable areas on fertile soils, the production method, the method of purchase or acquisition by the consumer, the environmental impact of the production method, the price
Regulating	Genetic resources	No studies were found for this ES based on the search terms.
	Nutrient retention	Denitrification rate, P retention level, soil type
	Biological control	Diversity of predator, non-native invasive species presence, potential for allelopathic effects, germination and seedling growth
	Carbon stock or sequestration, gas regulation and climate regulation	C content, stock and sequestration in soil and trees (above ground living biomass, diameter at breast height, height of trees, soil organic matter), N ₂ O production, hillslope broadleaf afforestation
	Disturbance regulation	Total storage capacity in the system and % of water body occupation in the floodplain, abundance and richness of animals and plants species, water quality indicators, and percentage of land use and cover
	Pollination	The area with potential pollinator nests, the distance between nests and fertile areas and number of visits of native pollinators to fertile areas
	Erosion control and sediment retention	Sediment and nutrient removal, sediment deposition, erosion rates, land use and land cover, soil parameters (temperature, moisture, increased bulk density, sand content, and pH), nutrient turnover from decomposition, net N mineralization rates, microbial C and N
	Water regulation	Physical indicators of water bodies (extent, width, water level), nutrient mass flow in water, flow watersheds, runoff volume, % land use and land cover, water quality indicators
Support	Nutrient cycling	Total organic carbon, total phosphorus, total nitrogen, denitrification potential, nitrogen in the microbial mass and nitrogen mineralization potential
	Soil formation	Soil parameters (% of sand/silt/clay, moisture, temperature, microbial activity, density and porosity)
	Nursery and refugia	Trees planted/area, tree cover, remnants of natural vegetation, diversity of bryophyte and, bird, mammals and fish population feature
Cultural	Recreation and aesthetic	Accessible areas with potential for visitation and noise pollution level

valuation of ES. The main methods applied were Benefit Transfer Method (BTM) and the contingent assessment method (CAM), also known as contingent valuation (MVC).

4. Discussion

4.1. Bibliometrics

The number of publications using at least one of the terms of riparian zones (riparian buffer or riparian vegetation or riparian zone or riparian areas or riparian forest) increased exponentially during the 21 years period and approximately 10% of them applied the ES evaluation (Figure 2a and b). We partially attribute this increase in studies regarding ES in riparian zones due to the influence of the Millennium Ecosystem Assessment (MEA, 2005). The ecosystem services evaluation makes it possible to assess the capacity of different ecosystems, that are part of the landscape, to provide ecosystem services and, above all, the trade-offs influenced by land use and management. That is, this approach

can assimilate the high complexity of natural and anthropogenic landscapes, offering the opportunity to understand how changes in environments affect the supply of ES. Thus, it can help us understand the ecological and social functions performed by natural ecosystems, strengthening the importance of conservation and rational use of the RZ.

In this study, we found that the countries that most published on RZ were: the United States, Brazil, Canada, China, Australia, and Germany. Rood et al. (2020) found a similar result: United States, Canada, China, Australia, Brazil, and England. Even though some countries, like Japan, have substantially published about riparian zones during our study period, they do not remain in the ranking in terms of the application of the ES approach. On the other hand, countries that traditionally have not published much on the theme of riparian zones, like South Africa, are standing out (in the number of publications) in applying the ES evaluation. We also highlight that the top 10 countries that have published the most about

Table 4. Main methods used for ecosystem service (ES) integration in riparian zones, the objective of analysis, and the associated references.

Methods	Objectives	Examples of references
Bayesian belief network (BBN)	Assessing SE provided by riparian zones	McVittie et al. (2015)
FyrisSKZ, working for water (WFW)	Assessing water quality	Turpie et al. (2008), Collettine et al. (2015)
Integrated valuation of ES and tradeoffs (InVEST)	Assessing the consequences of reducing riparian zones and ES pollination	Meehan et al. (2013), Garrastazú et al. (2015)
Soil and water assessment tool (SWAT)	Assessing sediment and water dynamics	Amatya et al. (2011), Palazón et al. (2014), Vigiak et al. (2016)
STELLA	Assessing interactions between water bodies and land use	Randhir & Ekness (2013)
CAESAR LISFLOOD (C-L)	Simulating the effect of in-channel obstructions on the distribution of erosion and deposition sediments	Walsh et al. (2020)
Riparian ecosystem management model (REMM)	Simulating ecological processes in riparian zones	Altier et al. (2002), Kim et al. (2007)
Copernicus land monitoring service (CLMS)	Assessing and monitoring green infrastructure in riparian zones	Piedelobo et al. (2019)
Land utilization capability indicator (LUCI)	Integrating land management decision support model applied to map areas providing ecosystem goods and services	Sharps et al. (2017)
Biotic integrity, Jaccard and Shanon-Weaver diversity indexes	Assessing the diversity and distribution of species and individuals in the riparian zones related to ES	Huh & Choi (2019), Burdon et al. (2020)
Riparian quality index (RQI), Stream visual assessment protocol (SVAP)	Assessing the quality of riparian zones	Bjorkland et al. (2001), Del Tánago & Jalón (2006), Warren II et al. (2015)
Normalized difference vegetation index (NDVI)	Assessing and mapping green infrastructure in riparian zones by remote sensing	Yang (2007), Fu et al. (2016)
Normalized difference water index (NDWI)	Assessing and mapping changes in water content in riparian zones by remote sensing	Pereira et al. (2018), Vanderhoof & Burt (2018)
Relative aggregated value of ecosystem services indexes (RAVES)	Prioritizing sites for ecological restoration based on ecosystem services	Comín et al. (2018)
Generalized additive models (GAM)	Describing the spatial pattern of pollination benefits using predictors	Lautenbach et al. (2011)
Relative pollination potential index (RPP)	Evaluating the benefits of pollination for crop yield	Zulian et al. (2013)
A new comprehensive ecosystem service index (CES)	Analyzing spatiotemporal changes and trade-offs to support the supply of multiple ecosystem services	Sun et al. (2018)
Water retention index	Planning land use to manage water resources	Vandecasteele et al. (2018)
Ecological limits of hydrologic alteration (ELOHA)	Developing environmental flow prescriptions for many streams and rivers in a user-defined geographic region or jurisdiction	Mackay et al. (2014)

riparian zones using the ES approach are spread through all the continents of the globe, which demonstrates the importance of the topic addressed at the global level.

4.2. Ecosystem services studies in riparian zones

Most of these studies were related to the Regulating category and aimed to analyze the ES

regarding the structure and functions of natural vegetation in riparian zones, ecological restoration processes, habitat maintenance, and aimed at improving water quality and climate regulation (Figure 2). There were some publications that performed a specific meta-analysis on this topic (e.g. Dybala et al, 2019). Currently, among the ES surveyed, the services that are being studied the

most are nutrient cycling, carbon sequestration, water supply, water regulation, and sediment retention (Angradi et al., 2016; De Sosa et al., 2018; Cole et al., 2020; Riis et al., 2020). Water purification and water stock were the most studied ecosystem functions (e.g. Yang et al., 2009; Lautenbach et al., 2011; Gutiérrez & Alonso, 2013), followed by the contribution to biodiversity (e.g. Gollan et al., 2013; Gray et al., 2014) and soil retention (e.g. Vigiak et al., 2016). The studies about nutrient flow have emphasized the erosion control and sediment retention ES, provided by riparian zones through sediment trapping and enhancement of the cohesion in root systems, providing channel stability and resistance to erosion (e.g. Vigiak et al., 2016).

4.3. Indicators of ecosystem services in riparian zones

The indicators used in the surveyed publications relate to different categories of ES, with emphasis on regulating (Table 3). The indicators that evaluated the cycling of nutrients were the most used based on the measurement of different values and attributes of the soil as well as the indicators used to assess the quantity and quality of water in riparian zones. These indicators are conventionally being used, regardless of the ES approach, over time.

Some indicators can measure more than one ES, mainly those related to water such as: disturbance regulation, water regulation, erosion control and sediment retention. But we chose to keep them separately in Table 3, as they are associated to the studies found from the search terms, which were related to each ES. While the studies that used the ES water regulation measured physical indicators related to the water flow, the disturbance regulation ES correlated these indicators with indicators of another nature such as those related to biodiversity or land use, aiming to identify disturbances in the riparian zone. Additionally, indicators related to pollination and cultural services appear in more recent publications, consistent with the ES approach, from MEA (2005).

It was observed either none or few studies (e.g. Sha et al., 2011; Kachenchart et al., 2012; Kaushal et al., 2014) focused on the dynamics and flows of SE. Ecosystem flows can be understood as the movement of substances, compounds, or nutrients in an ecosystem passing from one compartment to another (Jenkins, 2005). Identifying indicators to measure the dynamics of ecosystem services is a great challenge, but also an opportunity for futures researches on the theme.

4.4. Methodologies of integrated analysis of ecosystem service in riparian zones

The methodologies of integrated analysis of ES have not been widely used to evaluate riparian zones. Water supply and erosion control ES, often associated with nutrient cycling, stood out among the publications that applied an integrated approach of ES (methods presented in Table 4). The Soil and Water Assessment Tool (SWAT) was the model most frequently used to model the dynamics of rivers and the flow of sediments in them (e.g. Amatya et al., 2011; Arnold et al., 2012; Palazón et al., 2014; Arnold et al., 2014; Vigiak et al., 2016; Francesconi et al., 2016).

Other methods have also been used to assess the relationship between land use and water ES (Water retention index, STELLA, and CAESAR LISFLOOD), but also models and indexes to assess ES provision capacity or ecology integrity of riparian zones as a whole (e.g. Bayesian belief network, Riparian ecosystem management model, Riparian quality index, Stream visual assessment protocol, and A new comprehensive ecosystem service index) (Table 4).

Studies on the habitat and diversity function assessed the biodiversity of the riparian landscape, its ability to provide habitat and resources for different species (Cole et al., 2015; Cole et al., 2020), and its influence on the distribution of native and invasive species (Shaker et al., 2017). Models and indexes were also found to assess biotic integrity (Biotic integrity, Jaccard and Shanon-Weaver diversity indexes) and green infrastructure in riparian zones, priority areas for restoration like Copernicus land monitoring service, Relative aggregated value of ecosystem services index, and Normalized difference vegetation index. Most of the publications that studied the pollination ES (regulation) were associated with the services refuge (support) and food production (provision) (e.g. Lautenbach et al., 2011; Meehan et al., 2013). The main methods applied were: Relative pollination potential index, Generalized additive models, and Integrated valuation of ES and tradeoffs (InVEST). InVEST is a suite of models used to map and value the goods and services from nature that sustain and fulfill human life. It helps explore how changes in ecosystems can lead to changes in the flows of many different benefits to people (e.g. Nelson et al., 2009; Garrastazú, et al., 2015; Posner et al., 2016).

Methodologies of integrated analysis of ES are extremely important, as this is how they work in the ecosystem itself, interacting and providing joint

responses in a complex system. This approach seems to be more adequate to evaluate sustainability in riparian zones as a support for their management. Riparian zones play essential roles in water and landscape planning, in restoration of aquatic systems, and in catalyzing institutional and societal cooperation for these efforts (Naiman & Décamps, 1997). Only an integrated analysis of ES can effectively help decision-makers to make choices that mutually benefit ecosystems and populations (Muller et al., 2010).

4.5. Ecosystem services at landscape or spatial multiscale in riparian zones

Publication on landscape management focused on assessing the willingness to adopt riparian buffers, especially in the agricultural landscape, in addition to proposing ways of managing riparian zones to contribute to their greater ecosystem potential (e.g. Kenwick et al., 2009; Hefting et al., 2013; Angelstam & Lazdinis, 2017). Studies that focused on riparian corridors addressed the importance of ecological corridors in riparian zones to ensure the connectivity of habitats in fragmented landscapes (e.g. Clerici et al., 2014; Lee et al., 2014; Resasco et al., 2014). Regarding water quality, the studies focused on the evaluation of the riparian landscape's capacity to retain nutrients and sediments, in addition to its function of water filtration and purification that guarantee the quality of water for human supply (e.g. Nava-López et al., 2016).

Few studies have chosen the multiscale approach (e.g. Jones et al., 2010; Uriarte et al., 2011; Salo & Theobald, 2016; Biggs et al. 2019). This is probably due to riparian zones being considered only stretches of landscape. This fact can also be explained by the great complexity and the difficulty of carrying out studies in these areas, not only due to high costs, but also due to the difficulty of access to riparian zones (Busato et al., 2019). Landscape and spatial multiscale approaches are important in decision-making, especially in very heterogeneous and more complex landscapes (Power, 2010). Then, it is an opportunity to consider these aspects in the studies regarding ES. The ES of water supply and water regulation, for example, depend on runoff patterns in the landscape and are influenced by several biophysical factors.

4.6. Economic valuation of ecosystem services in riparian zones

Regarding economic valuation of ES, these methods estimate the monetary value of ES in

relation to other goods and services available in the economy, indicating a sustainable use of environmental market resources that makes viable use of environmental resources (e.g. Da Motta, 1997; Liu et al., 2010; Costanza et al., 2011). The main methods are BTM and MVC. BTM estimates the value of an ecosystem based on the results of pre-existing primary studies on different sites or policies (e.g. Liu et al., 2010; Rolfe et al., 2015). In the MVC, questionnaires are applied to individuals through direct consultation and simulation of contingent markets to reveal the willingness to pay (WTP) for the preservation or improvement of the ecosystem conditions, or the willingness to receive (WTR) some compensation for the loss of ecosystems conditions (e.g. Mueller, 2014; Garcia et al., 2015). Models for the economic valuation of ES, such as the Integrated Valuation of Ecosystem Services and Tradeoffs (InVEST), were also identified in the analyzed publications (e.g. Garrastazú et al., 2015). Studies aimed at supporting policies and markets, with a very applied focus related to carbon and other gases, assess to subsidize carbon credit markets and climate change adaptation (e.g. Matzek et al., 2014; Dittrich et al., 2018), as well as carrying out an economic valuation to support Payments for Ecosystem Services schemes (e.g. Lewis et al., 2017), have been increasingly frequent in the literature.

Valuing ES is important, as it shows understandably that investing in the maintenance of ecosystems can be more economical than later investing in actions for remediation, restoration and mitigation of the impacts arising from degradation. This approach is also key to subsidizing rising Payments for Ecosystem Services schemes around the world, encouraging conservation and greater provision of ES in RZ.

5. Conclusions

Despite the different definitions, functions, and high complexity, riparian zones have attracted the attention of scientists around the world who seek to understand their functioning and the provision of ecosystem services.

From 2000 to 2020, the most studied category of ecosystem services was Regulating, mostly carbon stock/sequestration, gas (GHG's) and climate regulation; biological control; pollination; erosion control and sediment retention; water regulation; nutrient retention, and sewage treatment. In the Provisioning category the ES most studied were water supply, food production, and raw material. The Support category included the ES of nutrient

cycling, soil formation, nursery, and refugia. In the Cultural category the ES studied the most were recreation and aesthetics.

The most applied indicators were those ones that evaluated the cycling of nutrients based on the measurement of soil and water attributes in riparian zones. Among the publications surveyed, it was noted that even with the use of riparian zone assessment and modeling indexes, few studies used methodologies of integrated analysis of ES. This fact calls attention to the need to carry out further studies that analyze ES in riparian zones using an integrated and multiscale approach because that is how the components of the ecosystem interact and provide joint responses capable of assisting in decision making.

Studies about ES in riparian zones using the landscape or spatial multiscale approach rarely appeared in the literature during the period studied. To assess the impacts of anthropogenic actions on ecosystem services, a spatial multiscale approach must be considered.

ES economic valuation was an approach rarely found in the publications surveyed in RZ, but it is very important to convince investors, decision-makers, and the society that it is better to invest in maintaining ES in riparian zones now than investing in remediation, restoration and mitigation of anthropogenic impacts later.

Finally, planning of water and landscape use, focusing on ES provision in RZ demand inter-institutional and social cooperation to strengthen management efforts, management, restoration and conservation of fauna, flora, water resources and ecosystem services in hydrographic basins, transforming conservation and sustainable use into a great opportunity for innovative development. Hence, connecting science and policy is essential to generate effective strategies for planning land use and monitoring ES's in natural and anthropized environments.

Acknowledgements

The authors would like to thank the infrastructure of Embrapa Soils and the Biosciences Institute of the Federal University of the State of Rio de Janeiro (UNIRIO).

References

Aguar Junior, T.R., & Parron, L.M., 2015. Indicadores de serviços ambientais hídricos e a contribuição da vegetação ripária para a qualidade de água. In: Parron, L.M., Garcia, J.R., Oliveira, E.B., Brown, G.G., &

Prado, R.B., eds. *Serviços ambientais em sistemas agrícolas e florestais do Bioma Mata Atlântica*. Brasília: Embrapa.

Altier, L.S., Lowrance, R., Williams, R.G., Inamdar, S.P., Bosh, D.D., Sheridan, J.M., Hubbard, R.K., & Thomas, D.L., 2002. *Riparian ecosystem management model: simulator for ecological processes in riparian zones*. USA: United States Department of Agriculture, Agricultural Research Service, Conservation Research Report, 46.

Amatya, D.M., Douglas-Mankin, K.R., Williams, T.M., Skaggs, R.W., & Nettles, J.E., 2011. Advances in forest hydrology: challenges and opportunities. *ASABE* 54(6), 2049-2056. <http://dx.doi.org/10.13031/2013.40672>.

Angelstam, P. & Lazdinis, M., 2017. Tall herb sites as a guide for planning, maintenance and engineering of riparian continuous forest cover. *Ecol. Eng.* 103(Pt B), 470-477. <https://doi.org/10.1016/j.ecoleng.2016.06.099>.

Angradi, T.R., Launspach, J.J., Bolgrien, D.W., Bellinger, B.J., Starry, M.A., Hoffman, J.C., Trebitz, A.S., Sierszen, M.E., & Hollenhorst, T.P., 2016. Mapping ecosystem service indicators in a Great Lakes estuarine Area of Concern. *J. Great Lakes Res.* 42(3), 717-727. <http://dx.doi.org/10.1016/j.jglr.2016.03.012>.

Arnold, J.G., Harmel, R.D., Johnson, M.-V.V., Bingner, R., Strickland, T.C., Walbridge, M., Santhi, C., Diluzio, M., & Wang, X., 2014. Impact of the agricultural research service watershed assessment studies on the conservation effects assessment project cropland national assessment. *J. Soil Water Conserv.* 69(5), 137A-144A. <http://dx.doi.org/10.2489/jswc.69.5.137A>.

Arnold, J.G., Moriasi, D.N., Gassman, P.W., Abbaspour, K.C., White, M.J., Srinivasan, R., Santhi, C., Harmel, R.D., Van Griensven, A., Van Liew, M.W., Kannan, N., & Jah, M.K., 2012. SWAT: model use, calibration, and validation. *ASABE* 55(41), 491-508. <https://doi.org/10.13031/2013.42256>.

Attanasio, C.M., Gandolfi, S., Zakia, M.J.B., Veniziani Junior, J.C.T., & Lima, W.P., 2012. A importância das áreas ripárias para a sustentabilidade hidrológica do uso da terra em microbacias hidrográficas. *Bragantia* 71(4), 493-501. <http://dx.doi.org/10.1590/S0006-87052013005000001>.

Biggs, T.W., Santiago, T.M.O., Sills, E., & Caviglia-Harris, J., 2019. The Brazilian Forest Code and riparian preservation areas: spatiotemporal analysis and implications for hydrological ecosystem services. *Reg. Environ. Change* 19(8), 2381-2394. <http://dx.doi.org/10.1007/s10113-019-01549-w>.

Bjorkland, R., Pringle, C.M., & Newton, B., 2001. A stream visual assessment protocol (SVAP) for riparian landowners. *Environ. Monit. Assess.* 68(2), 99-125. PMID:11411146. <http://dx.doi.org/10.1023/A:1010743124570>.

- Brauman, K.A., Daily, G.C., Duarte, T.K., & Mooney, H.A., 2007. The nature and value of ecosystem services: an overview highlighting hydrologic services. *Annu. Rev. Environ. Resour.*, 32, 67-98. <https://doi.org/10.1146/annurev.energy.32.031306.102758>.
- Bren, L.J., 1993. Riparian zone, stream, and floodplain issues: a review. *J. Hydrol. (Amst.)* 150(2-4), 277-299. [http://dx.doi.org/10.1016/0022-1694\(93\)90113-N](http://dx.doi.org/10.1016/0022-1694(93)90113-N).
- Burdon, F.J., Ramberg, E., Sargac, J., Forio, M.A.E., de Saeyer, N., Mutinova, P.T., Moe, T.F., Pavelescu, M.O., Dinu, V., Cazacu, C., Witing, F., Kupilas, B., Grandin, U., Volk, M., Rîșnoveanu, G., Goethals, P., Friberg, N., Johnson, R.K., & McKie, B.G., 2020. Assessing the benefits of forested riparian zones: a qualitative index of riparian integrity is positively associated with ecological status in European streams. *Water* 12(4), 1178. <http://dx.doi.org/10.3390/w12041178>.
- Busato, L., Boaga, J., Perri, M.T., Majone, B., Bellin, A., & Cassiani, G., 2019. Hydrogeophysical characterization and monitoring of the hyporheic and riparian zones: the Vermigliana Creek case study. *Sci. Total Environ.* 648, 1105-1120. PMID:30340257. <http://dx.doi.org/10.1016/j.scitotenv.2018.08.179>.
- Cao, Y., & Natuhara, Y., 2020. Effect of Urbanization on vegetation in riparian area: plant communities in artificial and semi-natural habitats. *Sustainability*, 12(1), 204. <https://doi.org/10.3390/su12010204>.
- Clerici, N., Paracchini, M.L., & Maes, J., 2014. Land-cover change dynamics and insights into ecosystem services in European stream riparian zones. *Ecohydrol. Hydrobiol.* 14(2), 107-120. <http://dx.doi.org/10.1016/j.ecohyd.2014.01.002>.
- Cole, L.J., Brocklehurst, S., Robertson, D., Harrison, W., & McCracken, D.I., 2015. Riparian buffer strips: their role in the conservation of insect pollinators in intensive grassland systems. *Agric. Ecosyst. Environ.* 211, 207-220. <http://dx.doi.org/10.1016/j.agee.2015.06.012>.
- Cole, L.J., Stockan, J., & Helliwell, R., 2020. Managing riparian buffer strips to optimize ecosystem services: a review. *Agric. Ecosyst. Environ.* 296, 106891. <http://dx.doi.org/10.1016/j.agee.2020.106891>.
- Collentine, D., Johnsson, H., Larsson, P., Markensten, H., & Persson, K., 2015. Designing cost efficient buffer zone programs: an application of the FyrisSKZ tool in a Swedish catchment. *Ambio* 44(Suppl. 2), S311- S 318. PMID:25681987. <http://dx.doi.org/10.1007/s13280-015-0627-y>.
- Comín, F.A., Beatriz, M., Ricardo, S., Felipe-Lucia, M.R., Jiménez, J.J., & Enrique, N., 2018. Prioritizing sites for ecological restoration based on ecosystem services. *J. Appl. Ecol.*, 55(3), 1155-1163. <https://doi.org/10.5061/dryad.8t5d4>.
- Costanza, R., D'Arge, R., De Groot, R., Farber, S., Grasso, M., Hannon, B., Limburg, K., Naeem, S., O'Neill, R.V., Paruelo, J., Raskin, R.G., Sutton, P., & Belt, M.V.D., 1997. The value of the world's ecosystem services and natural capital. *Nature*, 387, 253-260. <https://doi.org/10.1038/387253a0>.
- Costanza, R., De Groot, R., Braat, L., Kubiszewski, I., Fioramonti, L., Sutton, P., Farber, S., & Grasso, M., 2017. Twenty years of ecosystem services: how far have we come and how far do we still need to go? *Ecosyst. Serv.* 28, 1-16. <http://dx.doi.org/10.1016/j.ecoser.2017.09.008>.
- Costanza, R., Kubiszewski, I., Ervin, D., Bluffstone, R., Boyd, J.M., Brown, D., Chang, H., Dujon, V., Granek, E., Polasky, S., Shandas, V., & Yeakley, A., 2011. Valuing ecological systems and services. *F1000 Biol. Rep.* 3, 14. PMID:21876725. <http://dx.doi.org/10.3410/B3-14>.
- Da Motta, R.S., 1997. Manual para valoração econômica de recursos ambientais. Brasília: IPEA/MMA/PNUD/CNPQ.
- Dahl, T.E., 1990. Wetland losses in the United States: 1780's to 1980's [online]. Washington, DC: Department of The Interior, Fish and Wildlife Service. Retrieved in 2020, August 20, from <https://www.fws.gov/wetlands/Documents/Wetlands-Losses-in-the-United-States-1780s-to-1980s.pdf>
- De Groot, R.S., Wilson, M.A., & Boumans, R.M.J., 2002. A typology for the classification, description and valuation of ecosystem functions, goods and services. *Ecol. Econ.* 41(3), 393-408. [http://dx.doi.org/10.1016/S0921-8009\(02\)00089-7](http://dx.doi.org/10.1016/S0921-8009(02)00089-7).
- De Sosa, L.L., Glanville, H.C., & Marshal, M.R., Prysor Williams, A., & Jones, D.L., 2018. Quantifying the contribution of riparian soils to the provision of ecosystem services. *Sci. Total Environ.*, 624, 807-819. <https://doi.org/10.1016/j.scitotenv.2017.12.179>.
- Del Tánago, M.G., & Jalón, D.G., 2006. Índice RQI para la valoración de las riberas fluviales en el contexto de la directiva marco del agua. *Ing. Civ.*, 143, 97-108.
- Denyer, D., & Tranfield, D., 2009. Producing a systematic review. In: Buchanan, D. A., & Bryman, A., eds. *The SAGE handbook of organizational research methods*. London: Sage Publications Ltd.
- Dittrich, R., Ball, T., Wreford, A., Moran, D., & Spray, C.J., 2018. A cost-benefit analysis of afforestation as a climate change adaptation measure to reduce flood risk. *J. Flood. Risk Manage.* 12, e12482. <https://doi.org/10.1111/jfr3.12482>.
- Dufour, S., Rodríguez-González, P.M., & Laslier, M., 2019. Tracing the scientific trajectory of riparian vegetation studies: main topics approaches and needs in a globally changing world. *Sci. Total Environ.* 653, 1168-1185. PMID:30759557. <http://dx.doi.org/10.1016/j.scitotenv.2018.10.383>.
- Dybala, K.E., Matzek, V., Gardali, T., & Seavy, N.E., 2019. Carbon sequestration in riparian forests: a global synthesis and meta-analysis. *Glob. Chang.*

- Biol. 25(1), 57-67. PMID:30411449. <http://dx.doi.org/10.1111/gcb.14475>.
- Francesconi, W., Srinivasan, R., Pérez-Miñana, E., Willcock, S.P., & Quintero, M., 2016. Using the Soil and Water Assessment Tool (SWAT) to model ecosystem services: a systematic review. *J. Hydrol. (Amst.)* 535, 625-636. <http://dx.doi.org/10.1016/j.jhydrol.2016.01.034>.
- Fu, B., Li, Y., Wang, Y., Zhang, B., Yin, S., Zhu, H., & Xing, Z., 2016. Evaluation of ecosystem service value of riparian zone using land use data from 1986 to 2012. *Ecol. Indic.* 69, 873-881. <https://doi.org/10.1016/j.ecolind.2016.05.048>.
- Garcia, J.R., dos Reis, J.C., Moreira, J.M.M.A.P., & Ferronato, C., 2015. Considerações teórico-metodológicas sobre o processo de valoração dos recursos naturais. In: Parron, L. M., Garcia, J. R., Oliveira, E. B., Brown, G. G. & Prado, R. B., eds. *Serviços ambientais em sistemas agrícolas e florestais do Bioma Mata Atlântica*. Brasília: Embrapa.
- Garrastazú, M.C., Mendonça, S.D., Horokoski, T.T., Cardoso, D.J., Rosot, M.A.D., Nimmo, E.R., & Lacerda, A.E.B., 2015. Carbon sequestration and riparian zones: assessing the impacts of changing regulatory practices in Southern Brazil. *Land Use Policy* 42, 329-339. <http://dx.doi.org/10.1016/j.landusepol.2014.08.003>.
- Gollan, J.R., De Bruyn, L.L., Reid, N., & Wilkie, L., 2013. Monitoring the ecosystem service provided by dung beetles offers benefits over commonly used biodiversity metrics and a traditional trapping method. *J. Nat. Conserv.* 21(3), 183-188. <http://dx.doi.org/10.1016/j.jnc.2012.12.004>.
- Gray, C.L., Slade, E.M., Mann, D.J., & Lewis, O.T., 2014. Do riparian reserves support dung beetle biodiversity and ecosystem services in oil palm-dominated tropical landscapes? *Ecol. Evol.* 4(7), 1049-1060. PMID:24772282. <http://dx.doi.org/10.1002/ece3.1003>.
- Gregory, S.V., & Ashkenas, L., 1990. *Riparian Management Guide* [online]. Oregon: USDA Forest Service Pacific Northwest Region. Retrieved in 2020, August 20, from <https://andrewsforest.oregonstate.edu/sites/default/files/lter/pubs/pdf/pub1214.pdf>
- Gregory, S.V., Swanson, F.J., McKee, W.A., & Cummins, K., 1991. An ecosystem perspective of riparian zones. *Bioscience*. 41(8), 540-551. <https://doi.org/10.2307/1311607>.
- Grizzetti, B., Lanzanova, D., Liqueste, C., Reynaud, A., & Cardoso, A.C., 2016. Assessing water ecosystem services for water resource management. *Environ. Sci. Policy*. 61, 194-203. <https://doi.org/10.1016/j.envsci.2016.04.008>.
- Gutiérrez, M., & Alonso, M.L.S., 2013. Which are, what is their status and what can we expect from ecosystem services provided by Spanish rivers and riparian areas? *Biodivers. Conserv.* 22(11), 2469-2503. <http://dx.doi.org/10.1007/s10531-013-0532-2>.
- Hefting, M.M., Van Den Heuvel, R.N., & Verhoeven, J.T.A., 2013. Wetlands in agricultural landscapes for nitrogen attenuation and biodiversity enhancement: opportunities and limitations. *Ecol. Eng.* 56, 5-13. <http://dx.doi.org/10.1016/j.ecoleng.2012.05.001>.
- Huh, M.K., & Choi, B.K., 2019. Floristic characterization and biodiversity of riparian zones at the Gwangyang river, Korea. *Eur. J. Appl. Sci.*, 6(2), 24-38.
- Jenkins, C., 2005. *Nutrient flux assessment in the Port waterways*. Adelaide: Environmental Protection Agency.
- Jones, K.B., Slonecker, E.T., Nash, M.S., Neale, A.C., Wade, T.G., & Hamann, S., 2010. Riparian habitat changes across the continental United States (1972–2003) and potential implications for sustaining ecosystem services. *Landsc. Ecol.* 25(8), 1261-1275. <http://dx.doi.org/10.1007/s10980-010-9510-1>.
- Kachenchart, B., Jones, D.L., Gajasen, N., Edwards-Jones, G., & Limsakul, A., 2012. Seasonal nitrous oxide emissions from different land uses and their controlling factors in a tropical riparian ecosystem. *Agric. Ecosyst. Environ.* 158, 15-30. <http://dx.doi.org/10.1016/j.agee.2012.05.008>.
- Kaushal, S.S., Mayer, P.M., Vidon, P.G., Smith, R.M., Pennino, M.J., Newcomer, T.A., Duan, S., Welty, C., & Belt, K.T., 2014. Land use and climate variability amplify carbon, nutrient, and contaminant pulses: a review with management implications. *J. Am. Water Resour. Assoc.* 50(3), 585-614. <http://dx.doi.org/10.1111/jawr.12204>.
- Kenwick, R.A., Shammin, M.R., & Sullivan, W.C., 2009. Preferences for riparian buffers. *Landsc. Urban Plan.* 91(2), 88-96. <http://dx.doi.org/10.1016/j.landurbplan.2008.12.005>.
- Kim, I.J., Hutchinson, S.L., Hutchinson, J.M.S., & Young, C.B., 2007. Riparian ecosystem management model: sensitivity to soil, vegetation, and weather input parameters. *JAWRA*, 45(5), 1171-1182. <https://doi.org/10.1111/j.1752-1688.2007.00096.x>.
- Kobiyama, M., 2003. Conceitos de zona ripária e seus aspectos geobiohidrológicos. In: *Proceedings of I Seminário de Hidrologia Florestal: Zonas Ripárias, Brasil, Alfredo Wagner-SC:Brazil*. Retrieved in 2020, August 20, from <http://www.labhidro.ufsc.br/Artigos/Seminario%20Hidrologia%20Florestal%20%282003%29.pdf>
- Lautenbach, S., Kugel, C., Lausch, A., & Seppelt, R., 2011. Analysis of historic changes in regional ecosystem service provisioning using land use data. *Ecol. Indic.* 11(2), 676-687. <http://dx.doi.org/10.1016/j.ecolind.2010.09.007>.
- Lee, J.A., Chon, J., & Ahn, C., 2014. Planning landscape corridors in ecological infrastructure using least-cost path methods based on the value of ecosystem

- services. *Sustainability*. 6(11), 7564-7585. <http://dx.doi.org/10.3390/su6117564>.
- Lewis, S.E., Popp, J.S., English, L.A. & Odetola, O., 2017. Willingness to pay for riparian zones in an Ozark Watershed. *J. Water Resour. Plan. Manag.*, 143(5), 04017006. [https://doi.org/10.1061/\(ASCE\)WR.1943-5452.0000740](https://doi.org/10.1061/(ASCE)WR.1943-5452.0000740).
- Lind, L., Hasselquist, E.M., & Laudon, H., 2019. Towards ecologically functional riparian zones: a meta-analysis to develop guidelines for protecting ecosystem functions and biodiversity in agricultural landscapes. *J. Environ. Manage.*, 249, 109391. <https://doi.org/10.1016/j.jenvman.2019.109391>.
- Liu, S., Costanza, R., Farber, S., & Troy, A., 2010. Valuing ecosystem services: theory, practice, and the need for a transdisciplinary synthesis. *Ann. N. Y. Acad. Sci.* 1185(1), 54-78. PMID:20146762. <http://dx.doi.org/10.1111/j.1749-6632.2009.05167.x>.
- Mackay, S., Arthington, A.H., & James, C.S., 2014. Classification and comparison of natural and altered flow regimes to support an Australian trial of the Ecological Limits of Hydrologic Alteration framework. *Ecohydrology* 7(6), 1485-1507. <http://dx.doi.org/10.1002/eco.1473>.
- Matzek, V., Pulleston, C., & Gunn, J., 2014. Can carbon credits fund riparian forest restoration? *Restor. Ecol.* 61, 7-14. <https://doi.org/10.1111/rec.12153>.
- McKergow, L.A., Weaver, D.M., Prosser, I.P., Grayson, R.B., & Reed, A.E.G., 2003. Before and after riparian management: sediment and nutrient exports from a small agricultural catchment, Western Australia. *J. Hydrol. (Amst.)* 270(3-4), 253-272. [http://dx.doi.org/10.1016/S0022-1694\(02\)00286-X](http://dx.doi.org/10.1016/S0022-1694(02)00286-X).
- McVittie, A., Norton, L., Martin-Ortega, J., Siameti, I., Glenk, K., & Aalders, I., 2015. Operationalizing an ecosystem services-based approach using Bayesian Belief Networks: an application to riparian buffer strips. *Ecol. Econ.* 110, 15-27. <http://dx.doi.org/10.1016/j.ecolecon.2014.12.004>.
- Meehan, T.D., Gratton, C., Diehl, E., Hunt, N.D., Mooney, D.F., Ventura, S.J., Barham, B.L., & Jackson, R.D., 2013. Ecosystem service tradeoffs associated with switching from annual to perennial energy crops in riparian zones of the US Midwest. *PLoS One* 8(11), e80093. PMID:24223215. <http://dx.doi.org/10.1371/journal.pone.0080093>.
- Millenium Ecosystem Assessment – MEA, 2005. *Ecosystem and human well-being: synthesis*. Washington: Island Press.
- Mitsch, W.J., & Gosselink, J.G., 2000. The value of wetlands: importance of scale and landscape setting. *Ecol. Econ.* 35(1), 25-33. [http://dx.doi.org/10.1016/S0921-8009\(00\)00165-8](http://dx.doi.org/10.1016/S0921-8009(00)00165-8).
- Mueller, J.M., 2014. Estimating willingness to pay for watershed restoration in Flagstaff, Arizona using dichotomous-choice contingent valuation. *Forestry* 87(2), 327-333. <http://dx.doi.org/10.1093/forestry/cpt035>.
- Muller, F., De Groot, R., & Willemsen, L., 2010. Ecosystem services at the landscape scale: the need for integrative approaches. *Landsc. Online* 23, 1-11. <http://dx.doi.org/10.3097/LO.201023>.
- Naiman, R.J., & Décamps, H., 1997. The ecology of interfaces: riparian zones. *Annu. Rev. Ecol. Evol. Syst.* 28(1), 621-658. <http://dx.doi.org/10.1146/annurev.ecolsys.28.1.621>.
- Natural Resources Conservation Service – NRCS, 1997. Riparian Forest Buffer [online]. Seattle: USDA-NRCS-Watershed Science Institute. Retrieved in 2020, August 20, from <https://www.wcc.nrcs.usda.gov/ftpref/wntsc/strmRest/buffers/RiparianForestBufferJobSheet.pdf>
- Nava-López, M.Z., Diemont, S.A.W., Hall, M., & Ávila-Akerberg, V., 2016. Riparian buffer zone and whole watershed influences on river water quality: implications for ecosystem services near megacities. *Environ. Process* 3(2), 277-305. <http://dx.doi.org/10.1007/s40710-016-0145-3>.
- Nelson, E., Mendoza, G., Regetz, J., Polasky, S., Tallis, H., Cameron, R.D., Chan, K.M., Daily, G.C., Goldstein, J., Kareiva, P.M., Lonsdorf, E., Naidoo, R., Ricketts, T.H., & Shaw, M.R., 2009. Modeling multiple ecosystem services, biodiversity conservation, commodity production, and tradeoffs at landscape scales. *Front. Ecol. Environ.* 7(1), 4-11. <http://dx.doi.org/10.1890/080023>.
- Palazón, L., Gaspar, L., Latorre, B., Blake, W.H., & Navas, A., 2014. Evaluating the importance of surface soil contributions to reservoir sediment in alpine environments: a combined modelling and fingerprinting approach in the Posets-Maladeta Natural Park. *Solid Earth* 5(2), 963-978. <http://dx.doi.org/10.5194/se-5-963-2014>.
- Pereira, L.E., Amorim, G., Grigio, A.M., & Paranhos Filho, A.C., 2018. Análise comparativa entre métodos de índice de água por diferença normalizada (NDWI) em área úmida continental. *Anu. Inst. Geocienc.* 41(2), 654-662. https://doi.org/10.11137/2018_2_654_662.
- Piedelobo, L., Taramelli, A., Schiavon, E. & Valentini, E., 2019. Assessment of green infrastructure in riparian zones using copernicus programme. *Remote Sens.* 11(24), 2967. <https://doi.org/10.3390/rs11242967>.
- Posner, S., Verutes, G., Koh, I., Denu, D., & Ricketts, T., 2016. Global use of ecosystem service models. *Ecosyst. Serv.* 17, 131-141. <http://dx.doi.org/10.1016/j.ecoser.2015.12.003>.
- Power, A.G., 2010. Ecosystem services and agriculture: tradeoffs and synergies. *Philos Trans. R. Soc. Lond. B. Biol. Sci.* 365(1554), 2959-2971. PMID:20713396. <http://dx.doi.org/10.1098/rstb.2010.0143>.

- Randhir, T.O., & Ekness, P., 2013. Water quality change and habitat potential in riparian ecosystems. *Ecohydrol. Hydrobiol.* 13(3), 192-200. <http://dx.doi.org/10.1016/j.ecohyd.2013.09.001>.
- Resasco, J., Haddad, N.M., Orrock, J.L., Shoemaker, D., Brudvig, L.A., Damschen, E.I., Tewksbury, J.J., & Levey, D.J., 2014. Landscape corridors can increase invasion by an exotic species and reduce diversity of native species. *Ecology* 95(8), 2033-2039. PMID:25230454. <http://dx.doi.org/10.1890/14-0169.1>.
- Riis, T., Kelly-Quinn, M., Aguiar, F.C., Manolaki, P., Bruno, D., Bejarano, M.D., Clerici, N., Fernandes, M.R., Franco, J.C., Pettit, N., Portela, A.P., Tammeorg, O., Tammeorg, P., Rodriguez-González, P.M., & Dufour, S., 2020. Global overview of ecosystem services provided by riparian vegetation. *Bioscience* 70(6), 501-514. <http://dx.doi.org/10.1093/biosci/biaa041>.
- Rodrigues, R.R. 2000. Uma discussão nomenclatural das formações ciliares. In: Rodrigues, R.R., & Leitão Filho, H.F.L., eds. *Matas ciliares: conservação e recuperação*. São Paulo: EDUSP.
- Rolfe, J., Johnston, R.J., Rosenberger, R.S., & Brouwer, R., 2015. Introduction: benefit transfer of environmental and resource values. In: Johnston, R.J., Rolfe, J., Rosenberger, R.S., & Brouwer, R. *Benefit Transfer of Environmental and Resource Values: a guide for researchers and practitioners*. http://dx.doi.org/10.1007/978-94-017-9930-0_1.
- Rood, S. B., Scott, M.L., Dixon, M., González, E., Marks, C. O., Shafroth, P. B. & Volke, M., 2020. Ecological interfaces between land and flowing water: themes and trends in riparian research and management. *Wetlands*. 40, 1801-1811. <http://doi.org/10.1007/s13157-020-01392-4>.
- Salo, J.A., & Theobald, D.M., 2016. A Multi-scale, hierarchical model to map riparian zones: a multi-scale, hierarchical model to map riparian zones. *River Res. Appl.* 32(8), 1709-1720. <http://dx.doi.org/10.1002/rra.3019>.
- Salvador, J.L.G., 1987. Considerações sobre as matas ciliares e a implantação de reflorestamentos mistos nas margens de rios e reservatórios. São Paulo: CESP, Série Divulgação e Informação.
- SCImago, 2020. SJR – SCImago Journal & Country Rank [online]. Retrieved in 2020, August 29, from <https://www.scimagojr.com/index.php>
- Selles, I.M., 2001. Revitalização de rios – orientação técnica. Rio de Janeiro: SEMADS. Retrieved in 2020, August 29, from <https://www.docsity.com/pt/revitalizacao-de-rios-orientacao-tecnica-semads-gtz/4838204/>
- Sha, C., Mitsch, W.J., Mander, Ü., Lu, J., Batson, J., Zhang, L., & He, W., 2011. Methane emissions from freshwater riverine wetlands. *Ecol. Eng.* 37(1), 16-24. <http://dx.doi.org/10.1016/j.ecoleng.2010.07.022>.
- Shaker, R.R., Yakubov, A., Nick, S., Vennie-Vollrath, E., Ehlinger, T.J. & Forsythe, K.W., 2017. Predicting aquatic invasion in Adirondack lakes: a spatial analysis of lake and landscape characteristics. *Ecosphere*. 8(3) <http://dx.doi.org/10.1002/ecs2.1723>.
- Sharps, K., Masante, D., Thomas, A., Jackson, B., Redhead, J., May, L., Prosser, H., Cosby, B., Emmett, B., & Jones, L., 2017. Comparing strengths and weaknesses of three ecosystem services modeling tools in a diverse UK river catchment. *Sci. Total Environ.* 584-585, 118-130. PMID:28147292. <http://dx.doi.org/10.1016/j.scitotenv.2016.12.160>.
- Soman, S., Beyeler, S., Kraft, S.E., Thomas, D., & Winstanley, D., 2007. Ecosystem services from riparian areas: a brief summary of the literature [online]. USA: Illinois River Coordinator Council. Retrieved in 2020, August 20, from <http://www.verdevalley.org/RiparianAreas.pdf>
- Souza, M.C., 1999. Algumas considerações sobre vegetação ripária. *Cad. Biodivers.* 2(1), 4-10.
- Sun, X., Lu, Z., Li, F. & Crittenden, J.C., 2018. Urban expansion simulation and the spatio-temporal changes of ecosystem services, a case study in Atlanta Metropolitan area, USA. *Sci. Total Environ.*, 622-623, 974-987.
- Sweeney, B.W., & Newbold, J.D., 2014. Streamside forest buffer width needed to protect stream water quality, habitat, and organisms: a literature review. *J. Am. Water Resour. Assoc.* 50, 560-584. <https://doi.org/10.1111/jawr.12203>.
- Sweeney, B.W., Bott, T.L., Jackson, J.K., Kaplan, L.A., Newbold, J.D., Standley, L.J., Hession, W.C., & Horwitz, R.J., 2004. Riparian deforestation, stream narrowing, and loss of stream ecosystem services. *Proc. Natl. Acad. Sci. USA* 101(39), 14132-14137. PMID:15381768. <http://dx.doi.org/10.1073/pnas.0405895101>.
- Tabacchi, E., Lambs, L., Guilloy, H., Planty-Tabacchi, A.M., Muller, E., & Décamps, H., 2000. Impacts of riparian vegetation on hydrological processes. *Hydrol. Processes* 14, 2959-2976. [http://dx.doi.org/10.1002/1099-1085\(200011/12\)14:16/17<2959::AID-HYP129>3.0.CO;2-B](http://dx.doi.org/10.1002/1099-1085(200011/12)14:16/17<2959::AID-HYP129>3.0.CO;2-B).
- Tansley, A.G., 1935. The use and abuse of vegetational concepts and terms. *Ecology* 16(3), 284-307. <http://dx.doi.org/10.2307/1930070>.
- Tiegs, S.D., Costello, D.M., Isken, M.W., Woodward, G., Mcintyre, P.B., Gessner, M.O., Chauvet, E., Griffiths, N.A., Flecker, A., & Zwart, J.A., 2019. Global patterns and drivers of ecosystem functioning in rivers and riparian zones. *Science*. 5(1), 1-8. <https://doi.org/10.1126/sciadv.aav0486>.

- Torres, R.B., Matthes, L.A.F., Rodrigues, R.R., & Leitão Filho, H.F., 1992. Lista de espécies florestais nativas para plantio em áreas de brejo. *O Agrônomo*. (Online), 44(1-3), 1-2. Retrieved in 2020, August 20, from http://www.fundacaofia.com.br/gdusm/lista_florestas_brejo.pdf
- Turpie, J.K., Marais, C., & Blignaut, J.N., 2008. The working for water programme: evolution of a payments for ecosystem services mechanism that addresses both poverty and ecosystem service delivery in South Africa. *Ecol. Econ.* 65(4), 788-798. <http://dx.doi.org/10.1016/j.ecolecon.2007.12.024>.
- Uriarte, M., Yackulic, C.B., Lim, Y., & Arce-Nazario, J.A., 2011. Influence of land use on water quality in a tropical landscape: a multi-scale analysis. *Landsc. Ecol.* 26(8), 1151-1164. PMID:26146455. <http://dx.doi.org/10.1007/s10980-011-9642-y>.
- Vandecasteele, I., Marí i Rivero, I., Baranzelli, C., Becker, W., Dreoni, I., Lavalle, C., & Batelaan, O., 2018. The Water Retention Index: using land use planning to manage water resources in Europe. *Sustain. Dev. (Bradford)* 26(2), 122-131. <http://dx.doi.org/10.1002/sd.1723>.
- Vanderhoof, M. & Burt, C., 2018. Applying high resolution imagery to evaluate restoration induced changes in stream condition, Missouri River Headwaters Basin, Montana. *Remote Sens.* 10(6), 913. <https://doi.org/10.3390/rs10060913>.
- Vigiak, O., Malagó, A., Bouraoui, F., Grizzetti, B., Weissteiner, C.J., & Pastori, M., 2016. Impact of current riparian land on sediment retention in the Danube River Basin. *Sustain. Water Qual. Ecol.* 8, 30-49. <http://dx.doi.org/10.1016/j.swaqe.2016.08.001>.
- Vitali, P.D.L., Zakia, M.J.B., & Durigan, G., 2009. Considerações sobre a legislação correlata à zonatampão de unidades de conservação no Brasil. *Ambiente Soc.* 12(1), 67-82. <http://dx.doi.org/10.1590/S1414-753X2009000100006>.
- Walsh, P., Jakeman, A., & Thompson, C., 2020. Modelling headwater channel response and suspended sediment yield to in-channel large wood using the Caesar-Lisflood landscape evolution model. *Geomorphology* 363, 107209. <http://dx.doi.org/10.1016/j.geomorph.2020.107209>.
- Warren II, R.J., Potts, D.L., & Frothingham, K.M., 2015. Stream structural limitations on invasive communities in urban riparian areas. *Invasive Plant Sci. Manag.* 8(3), 353-362. <http://dx.doi.org/10.1614/IPSM-D-14-00081.1>.
- Webb, A.A., & Erskine, W.D., 2003. A practical scientific approach to riparian vegetation rehabilitation in Australia. *J. Environ. Manage.* 68(4), 329-341. PMID:12877867. [http://dx.doi.org/10.1016/S0301-4797\(03\)00071-9](http://dx.doi.org/10.1016/S0301-4797(03)00071-9).
- Yang, L.R., Chen, L.D., & Sun, R.H., 2009. River ecosystems and their self-purification capability: Research status and challenges. *Shengtai Xuebao/Acta Ecologica Sinica* 29, 5066-5075.
- Yang, X., 2007. Integrated use of remote sensing and geographic information systems in riparian vegetation delineation and mapping. *Remote Sens.* 28(2), 353-370. <http://dx.doi.org/10.1080/01431160600726763>.
- Zulian, G., Maes, J., & Paracchini, M.L., 2013. Linking land cover data and crop yields for mapping and assessment of pollination services in Europe. *Land (Basel)* 2(3), 472-492. <http://dx.doi.org/10.3390/land2030472>.

Received: 03 March 2022

Accepted: 21 June 2022

Associated Editor: André Andrian Padial.